

# **ORF522 – Linear and Nonlinear Optimization**

## **4. The simplex method**


# Ed Forum

- Notebooks on GitHub: <https://github.com/ORF522/companion>
- Office hours change:  
Prof. Stellato: Thu 3:30pm-5:30pm  
Scander Mustapha: Mon: 1:30pm-3:30pm
- 10% Participation. The note should **summarize what you learned** in the last lecture, and **highlight the concepts that were most confusing** or that you would like to review. A note will receive full credit if: it is **submitted before the beginning of next lecture**, it is **related to the content** of the lecture, and it is **understandable** and coherent.
- Question: connection between geometry and standard form?  
Yes, they are equivalent (more in the next slides)

# Recap

# Equivalence Theorem

Given a nonempty polyhedron  $P = \{x \mid Ax \leq b\}$



Let  $x \in P$

$x$  is a **vertex**  $\iff x$  is an **extreme point**  $\iff x$  is a **basic feasible solution**

# Basic feasible solution

$$P = \{x \mid a_i^T x \leq b_i, \quad i = 1, \dots, m\}$$

**Active constraints at  $\bar{x}$**

$$\mathcal{I}(\bar{x}) = \{i \in \{1, \dots, m\} \mid a_i^T \bar{x} = b_i\}$$


Index of all the constraints  
satisfied as **equality**

**Basic solution  $\bar{x}$**

- $\{a_i \mid i \in \mathcal{I}(\bar{x})\}$  has  $n$  linearly independent vectors

**Basic feasible solution  $\bar{x}$**

- $\bar{x} \in P$
- $\{a_i \mid i \in \mathcal{I}(\bar{x})\}$  has  $n$  linearly independent vectors



# Standard form polyhedra

## Definition

### Standard form LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

## Assumption


$A \in \mathbf{R}^{m \times n}$  has full row rank  $m \leq n$

## Interpretation

$P$  lives in  $(n - m)$ -dimensional subspace

## Standard form polyhedron

$$P = \{x \mid Ax = b, x \geq 0\}$$



# Basic solutions

## Standard form polyhedra

$$P = \{x \mid Ax = b, x \geq 0\} \quad \text{with} \quad A \in \mathbf{R}^{m \times n} \text{ has full row rank } m \leq n$$

$x$  is a **basic solution** if and only if

- $Ax = b$
- There exist indices  $B(1), \dots, B(m)$  such that
  - columns  $A_{B(1)}, \dots, A_{B(m)}$  are linearly independent
  - $x_i = 0$  for  $i \neq B(1), \dots, B(m)$

$x$  is a **basic feasible solution** if  $x$  is a **basic solution** and  $x \geq 0$

# From geometry to standard form

$$\begin{array}{ll}
 \text{minimize} & c^T x \\
 \text{subject to} & Ax \leq b
 \end{array}
 \longrightarrow
 \begin{array}{ll}
 \text{minimize} & c^T (x^+ - x^-) \\
 \text{subject to} & \begin{bmatrix} A & -A & I \end{bmatrix} \begin{bmatrix} x^+ \\ x^- \\ s \end{bmatrix} = b \\
 & (x^+, x^-, s) \geq 0
 \end{array}
 \longrightarrow
 \begin{array}{ll}
 \text{minimize} & \tilde{c}^T \tilde{x} \\
 \text{subject to} & \tilde{A} \tilde{x} = b \\
 & \tilde{x} \geq 0
 \end{array}$$

Variables:  $\tilde{n} = 2n + m$

(Equality) constraints:  $\tilde{m} = m \implies$  **active**

Formal proof at  
Theorem 2.4 LO book

For a **basic solution**  $\longrightarrow$  We need  $\tilde{n} - \tilde{m} = 2n$   
active inequalities  $\Rightarrow \tilde{x}_i = 0$  (non basic)

Which corresponds to  $m$  inequalities inactive  $\Rightarrow \tilde{x}_i > 0$  (basic)



# Constructing basic solution

1. Choose any  $m$  independent columns of  $A$ :  $A_{B(1)}, \dots, A_{B(m)}$
2. Let  $x_i = 0$  for all  $i \neq B(1), \dots, B(m)$
3. Solve  $Ax = b$  for the remaining  $x_{B(1)}, \dots, x_{B(m)}$

$$\begin{array}{c} \text{Basis} \\ \text{matrix} \end{array} \quad \begin{array}{c} \text{Basis columns} \end{array} \quad \begin{array}{c} \text{Basic variables} \end{array}$$

$$A_B = \left[ \begin{array}{c|c|c|c} | & | & & | \\ A_{B(1)} & A_{B(2)} & \dots & A_{B(m)} \\ | & | & & | \end{array} \right], \quad x_B = \begin{bmatrix} x_{B(1)} \\ \vdots \\ x_{B(m)} \end{bmatrix} \longrightarrow \text{Solve } A_B x_B = b$$


If  $x_B \geq 0$ , then  $x$  is a **basic feasible solution**

# Optimality of extreme points

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

- If
- $P$  has at least one extreme point
  - There exists an optimal solution  $x^*$


Then, there exists an optimal solution which is an **extreme point** of  $P$



We only need to search between **extreme points**

# Conceptual algorithm

- Start at corner
- Visit neighboring corner that improves the objective



# Today's agenda

**Readings: [Chapter 3, LO]**

## **Simplex method**

- Iterate between neighboring basic solutions
- Optimality conditions
- Simplex iterations

# The simplex method

## Top 10 algorithms of the 20th century

1946: Metropolis algorithm

**1947: Simplex method** —————→

1950: Krylov subspace method

1951: The decompositional approach to matrix computations

1957: The Fortran optimizing compiler

1959: QR algorithm


1962: Quicksort

1965: Fast Fourier transform

1977: Integer relation detection

1987: Fast multipole method


George Dantzig



**Neighboring basic solutions**

# Neighboring solutions

Two basic solutions are **neighboring** if their basic indices differ by exactly one variable



## Example

$$\begin{matrix} & & A & & \\ \begin{bmatrix} 1 & -1 & 0 & 3 & -2 \\ 2 & 0 & -1 & -1 & 0 \\ 0 & 2 & 4 & -1 & 4 \end{bmatrix} & \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} & = & \begin{matrix} b \\ \begin{bmatrix} -5 \\ -1 \\ 14 \end{bmatrix} \end{matrix}$$

$$B = \{1, 3, 5\} \quad x_2 = x_4 = 0$$

$$A_B x_B = b \longrightarrow x_B = \begin{bmatrix} x_1 \\ x_3 \\ x_5 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 2.5 \end{bmatrix}$$

$$\bar{B} = \{1, 3, 4\} \quad y_2 = y_5 = 0$$

$$A_{\bar{B}} y_{\bar{B}} = b \longrightarrow y_{\bar{B}} = \begin{bmatrix} y_1 \\ y_3 \\ y_4 \end{bmatrix} = \begin{bmatrix} 0.1 \\ 3.0 \\ -1.7 \end{bmatrix}^{15}$$

# Feasible directions

## Conditions


$$P = \{x \mid Ax = b, x \geq 0\}$$

Given a basis matrix  $A_B = \begin{bmatrix} A_{B(1)} & \dots & A_{B(m)} \end{bmatrix}$

we have basic feasible solution  $x$ :

- $x_B$  solves  $A_B x_B = b$
- $x_i = 0, \forall i \neq B(1), \dots, B(m)$

Let  $x \in P$ , a vector  $d$  is a **feasible direction** at  $x$   
if  $\exists \theta > 0$  for which  $x + \theta d \in P$



### Feasible direction $d$

- $A(x + \theta d) = b \implies Ad = 0$
- $x + \theta d \geq 0$



# Feasible directions

## Computation

### Feasible direction $d$

- $A(x + \theta d) = b \implies Ad = 0$
- $x + \theta d \geq 0$

### Nonbasic indices

- $d_j = 1 \longrightarrow$  **Basic direction**
- $d_k = 0, \forall k \notin \{j, B(1), \dots, B(m)\}$

### Basic indices

$$Ad = 0 = \sum_{i=1}^n A_i d_i = A_B d_B + A_j = 0 \implies d_B = -A_B^{-1} A_j$$

### Non-negativity (non-degenerate assumption)

- Non-basic variables:  $x_i = 0$ . Nonnegative direction  $d_i \geq 0$
- Basic variables:  $x_B > 0$ . Therefore  $\exists \theta > 0$  such that  $x_B + \theta d_B \geq 0$

# Feasible directions

## Example


$$P = \{x \mid x_1 + x_2 + x_3 = 2, \quad x \geq 0\}$$

$$x = (2, 0, 0) \quad B = \{1\}$$

**Nonbasic index**  $j = 3 \longrightarrow d = (-1, 0, 1)$

$$d_j = 1$$

$$A_B d_B = -A_j \quad \Rightarrow \quad d_B = -1$$



# How does the cost change?

**Cost improvement**

$$c^T(x + \theta d) - c^T x = \theta c^T d$$

**New cost**

**Old cost**

We call  $\bar{c}_j$  the **reduced cost** of  
(introducing) variable  $x_j$  in the basis

$$\bar{c}_j = c^T d = \sum_{i=1}^n c_i d_i = c_j + c_B^T d_B = c_j - c_B^T A_B^{-1} A_j$$

# Reduced costs

## Interpretation

Change in objective/marginal cost of adding  $x_j$  to the basis

$$\bar{c}_j = c_j - c_B^T A_B^{-1} A_j$$

Cost per-unit increase  
of variable  $x_j$

Cost to change other variables  
compensating for  $x_j$   
to enforce  $Ax = b$

- $\bar{c}_j > 0$ : adding  $x_j$  will increase the objective (bad)
- $\bar{c}_j < 0$ : adding  $x_j$  will decrease the objective (good)

## Reduced costs for basic variables is 0

$$\begin{aligned}\bar{c}_{B(i)} &= c_{B(i)} - c_B^T A_B^{-1} A_{B(i)} = c_{B(i)} - c_B^T (A_B^{-1} A_B) e_i \\ &= c_{B(i)} - c_B^T e_i = c_{B(i)} - c_{B(i)} = 0\end{aligned}$$

# Vector of reduced costs

## Reduced costs

$$\bar{c}_j = c_j - c_B^T A_B^{-1} A_j$$

## Full vector in one shot?

$$\bar{c} = (\bar{c}_1, \dots, \bar{c}_n)$$

Isolate basis  $B$ -related components  $p$   
(they are the same across  $j$ )

$$\bar{c}_j = c_j - A_j^T (A_B^{-1})^T c_B = c_j - A_j^T p$$

Obtain  $p$  by solving linear system

$$p = (A_B^{-1})^T c_B \quad \Rightarrow \quad A_B^T p = c_B$$

Note:  $(M^{-1})^T = (M^T)^{-1}$   
for any square invertible  $M$

## Computing reduced cost vector

1. Solve  $A_B^T p = c_B$
2.  $\bar{c} = c - A^T p$

# Optimality conditions

# Optimality conditions

## Theorem

Let  $x$  be a basic feasible solution associated with basis matrix  $A_B$

Let  $\bar{c}$  be the vector of reduced costs.

If  $\bar{c} \geq 0$ , then  $x$  is **optimal**

## Remark

This is a **stopping criterion** for the simplex algorithm.

If the **neighboring solutions** do not improve the cost, we are done (because of convexity).

# Optimality conditions

## Proof

For a basic feasible solution  $x$  with basis  $B$  the reduced costs are  $\bar{c} \geq 0$ .

Consider any feasible solution  $y$  and define  $d = y - x$

Since  $x$  and  $y$  are feasible, then  $Ax = Ay = b$  and  $Ad = 0$

$$Ad = A_B d_B + \sum_{i \in N} A_i d_i = 0 \quad \Rightarrow \quad d_B = - \sum_{i \in N} A_B^{-1} A_i d_i$$


$N$  are the  
nonbasic indices

The change in objective is

$$c^T d = c_B^T d_B + \sum_{i \in N} c_i d_i = \sum_{i \in N} (c_i - c_B^T A_B^{-1} A_i) d_i = \sum_{i \in N} \bar{c}_i d_i$$

Since  $y \geq 0$  and  $x_i = 0, i \in N$ , then  $d_i = y_i - x_i \geq 0, i \in N$

$$c^T d = c^T (y - x) \geq 0 \quad \Rightarrow \quad c^T y \geq c^T x.$$





# Simplex iterations

# Stepsize

What happens if some  $\bar{c}_j < 0$ ?

We can decrease the cost by bringing  $x_j$  into the basis

**How far can we go?**

$$\theta^* = \max\{\theta \mid \theta \geq 0 \text{ and } x + \theta d \geq 0\} \quad d \text{ is the } j\text{-th basic direction}$$

**Unbounded**

If  $d \geq 0$ , then  $\theta^* = \infty$ . The LP is unbounded.

**Bounded**

If  $d_i < 0$  for some  $i$ , then

$$\theta^* = \min_{\{i \mid d_i < 0\}} \left( -\frac{x_i}{d_i} \right) = \min_{\{i \in B \mid d_i < 0\}} \left( -\frac{x_i}{d_i} \right)$$

(Since  $d_i \geq 0$ ,  $i \notin B$ )

# Moving to a new basis

## Next feasible solution

$$x + \theta^* d$$

Let  $B(\ell) \in \{B(1), \dots, B(m)\}$  be the index such that  $\theta^* = -\frac{x_{B(\ell)}}{d_{B(\ell)}}$ . Then,

$$x_{B(\ell)} + \theta^* d_{B(\ell)} = 0$$

## New solution

- $x_{B(\ell)}$  becomes 0 (exits)
- $x_j$  becomes  $\theta^*$  (enters)

## New basis

$$A_{\bar{B}} = \begin{bmatrix} A_{B(1)} & \dots & A_{B(\ell-1)} & A_j & A_{B(\ell+1)} & \dots & A_{B(m)} \end{bmatrix}$$

# An iteration of the simplex method

## First part

We start with

- a basic feasible solution  $x$
- a basis matrix  $A_B = \begin{bmatrix} A_{B(1)} & \dots, A_{B(m)} \end{bmatrix}$

1. Compute the reduced costs  $\bar{c}$

- Solve  $A_B^T p = c_B$
- $\bar{c} = c - A^T p$

2. If  $\bar{c} \geq 0$ ,  $x$  **optimal. break**

3. Choose  $j$  such that  $\bar{c}_j < 0$

# An iteration of the simplex method

## Second part

4. Compute search direction  $d$  with  $d_j = 1$  and  $A_B d_B = -A_j$
5. If  $d_B \geq 0$ , the problem is **unbounded** and the optimal value is  $-\infty$ . **break**
6. Compute step length  $\theta^* = \min_{\{i \in B \mid d_i < 0\}} \left( -\frac{x_i}{d_i} \right)$
7. Define  $y$  such that  $y = x + \theta^* d$
8. Get new basis  $\bar{B}$  ( $i$  exits and  $j$  enters)

# Example

$$P = \{x \mid x_1 + x_2 + x_3 = 2, \quad x \geq 0\}$$


$$x = (2, 0, 0) \quad B = \{1\}$$

$$\text{Basic index } j = 3 \longrightarrow d = (-1, 0, 1) \\ d_j = 1$$

$$A_B d_B = -A_j \quad \Rightarrow \quad d_B = -1$$

$$\text{Stepsize } \theta^* = -\frac{x_1}{d_1} = 2$$

$$\text{New solution } y = x + \theta^* d = (0, 0, 2) \quad \bar{B} = \{3\}$$



# Finite convergence

**Assume that**

- $P = \{x \mid Ax = b, x \geq 0\}$  not empty
- Every basic feasible solution **non degenerate**

**Then**

- The simplex method **terminates after a finite number of iterations**
- At termination we either have one of the following
  - an **optimal basis**  $B$
  - a **direction**  $d$  such that  $Ad = 0$ ,  $d \geq 0$ ,  $c^T d < 0$  and the optimal cost is  $-\infty$

# Finite convergence

## Proof sketch

At each iteration the algorithm improves


- by a **positive** amount  $\theta^*$
- along the **direction**  $d$  such that  $c^T d < 0$

Therefore

- The cost strictly decreases
- No basic feasible solution can be visited twice

Since there is a **finite number of basic feasible solutions**

The algorithm **must eventually terminate**





# The simplex method

Today, we learned to:

- **Iterate** between basic feasible solutions
- **Verify** optimality and unboundedness conditions
- **Apply** a single iteration of the simplex method
- **Prove** finite convergence of the simplex method in the non-degenerate case

# Next lecture

- Finding initial basic feasible solution
- Degeneracy
- Complexity