

ORF522 – Linear and Nonlinear Optimization

12. Introduction to nonlinear optimization

Homogeneous self-dual embedding

Optimality conditions

Primal

$$\begin{aligned} &\text{minimize} && c^T x \\ &\text{subject to} && Ax + s = b \\ &&& s \geq 0 \end{aligned}$$

Dual

$$\begin{aligned} &\text{maximize} && -b^T y \\ &\text{subject to} && A^T y + c = 0 \\ &&& y \geq 0 \end{aligned}$$

Optimality conditions

$$\begin{bmatrix} 0 \\ s \\ 0 \end{bmatrix} = \begin{bmatrix} 0 & A^T \\ -A & 0 \\ c^T & b^T \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} c \\ b \\ 0 \end{bmatrix}$$

$$s, y \geq 0$$

Any (x^*, s^*, y^*) satisfying these conditions is **optimal**

What happens if the problem is infeasible?

How do you detect infeasibility/unboundedness?

Primal

$$\begin{aligned} &\text{minimize} && c^T x \\ &\text{subject to} && Ax + s = b \\ &&& s \geq 0 \end{aligned}$$

Dual

$$\begin{aligned} &\text{maximize} && -b^T y \\ &\text{subject to} && A^T y + c = 0 \\ &&& y \geq 0 \end{aligned}$$

Alternatives (Farkas lemma) Write feasibility problem and dualize...

- **primal feasible:** $Ax + s = b, \quad s \geq 0$
- **primal infeasible:** $A^T y = 0, \quad b^T y < 0, \quad y \geq 0$ (primal infeasibility certificate)
- **dual feasible:** $A^T y + c = 0, \quad y \geq 0$
- **dual infeasible:** $Ax \leq 0, \quad c^T x < 0$ (dual infeasibility certificate)

The homogeneous self-dual embedding

Derivation

Introduce two new variables $\kappa, \tau \geq 0$

Homogeneous self-dual embedding

$$\begin{bmatrix} 0 \\ s \\ \kappa \end{bmatrix} = \begin{bmatrix} 0 & A^T & c \\ -A & 0 & b \\ -c^T & -b^T & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ \tau \end{bmatrix}$$

$$s, y, \kappa, \tau \geq 0$$



$$Qu = v$$

$$u, v \geq 0$$

$$Q = \begin{bmatrix} 0 & A^T & c \\ -A & 0 & b \\ -c^T & -b^T & 0 \end{bmatrix}$$

$$u = (x, y, \tau)$$

$$v = (0, s, \kappa)$$

The homogeneous self-dual embedding

Properties

$$Qu = v$$

$$u, v \geq 0$$

$$Q = \begin{bmatrix} 0 & A^T & c \\ -A & 0 & b \\ -c^T & -b^T & 0 \end{bmatrix}$$

$$u = (x, y, \tau)$$

$$v = (0, s, \kappa)$$

Matrix

- Q is skew-symmetric: $Q^T = -Q \Rightarrow u^T Qu = 0$

- $u \perp v$ **proof** $Qu - v = 0 \Rightarrow u^T Qu - u^T v = 0 \Rightarrow u^T v = 0$ ■

Homogeneous

(u, v) satisfy $Qu = v, (v, u) \geq 0 \Rightarrow \alpha(u, v)$ with $\alpha \geq 0$ feasible

Always feasible

$\alpha = 0 \Rightarrow (0, 0)$ is feasible

The homogeneous self-dual embedding

Outcomes

Find x, s, y, κ, τ such that

$$\begin{bmatrix} 0 \\ s \\ \kappa \end{bmatrix} = \begin{bmatrix} 0 & A^T & c \\ -A & 0 & b \\ -c^T & -b^T & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ \tau \end{bmatrix}$$
$$s, y, \kappa, \tau \geq 0$$

Note. By strict complementarity, we can ensure $\kappa + \tau > 0$

Case 1: feasibility

$\tau > 0, \kappa = 0$ define $(\hat{x}, \hat{s}, \hat{y}) = (x^*/\tau, s^*/\tau, y^*/\tau)$

$$\begin{aligned} 0 &= A^T \hat{y} + c & \hat{s} &\geq 0, & \hat{y} &\geq 0, & \hat{s}^T \hat{y} &= 0 \\ \hat{s} &= -A\hat{x} + b \end{aligned}$$

→ $(\hat{x}, \hat{s}, \hat{y})$ is a **solution** to the original problem

The homogeneous self-dual embedding

Outcomes

$$\text{Find } x, s, y, \kappa, \tau \text{ such that } \begin{bmatrix} 0 \\ s \\ \kappa \end{bmatrix} = \begin{bmatrix} 0 & A^T & c \\ -A & 0 & b \\ -c^T & -b^T & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ \tau \end{bmatrix}$$
$$s, y, \kappa, \tau \geq 0$$

Case 2: infeasibility

$\tau = 0, \kappa > 0 \longrightarrow c^T x + b^T y < 0$ (**impossible**). Must have infeasibility

If $b^T y < 0$ then $\hat{y} = y / (-b^T y)$ is a **certificate of primal infeasibility**

$$A^T \hat{y} = 0, \quad b^T \hat{y} = -1 < 0, \quad \hat{y} \geq 0$$

If $c^T x < 0$ then $\hat{x} = x / (-c^T x)$ is a **certificate of dual infeasibility**

$$A \hat{x} \leq 0, \quad c^T \hat{x} = -1 < 0$$

Interior-point method for homogeneous self-dual embedding

Linear complementarity problem

$$Qu = v$$

$$u^T v = 0$$

$$u, v \geq 0$$

Equations

$$h(u, v) = \begin{bmatrix} Qu - v \\ UV\mathbf{1} \end{bmatrix} = 0$$

$$u, v \geq 0$$

Directions

$$\begin{bmatrix} Q & -I \\ V & U \end{bmatrix} \begin{bmatrix} \Delta u \\ \Delta v \end{bmatrix} = \begin{bmatrix} -r_e \\ -UV\mathbf{1} + \sigma\mu\mathbf{1} \end{bmatrix}$$

$$r_e = Qu - v$$

$$\mu = (u^T v) / d$$

Line search to enforce $u, v > 0$

$$(u, v) \leftarrow (u, v) + \alpha(\Delta u, \Delta v)$$

Today's lecture

[Chapter 2-4 and 6, CO] [Chapter A and B, FCA]

- Nonlinear optimization
- Examples
- Convex analysis review
- Convex optimization

What if the problem is no longer linear?

Nonlinear optimization

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & g_i(x) \leq 0, \quad i = 1, \dots, m \end{array}$$

$x = (x_1, \dots, x_n)$ Variables

$f : \mathbf{R}^n \rightarrow \mathbf{R}$ Nonlinear objective function


$g_i : \mathbf{R}^n \rightarrow \mathbf{R}$ Nonlinear constraints functions

Feasible set

$$C = \{x \mid g_i(x) \leq 0, \quad i = 1, \dots, m\}$$

Small example

$$\begin{aligned} &\text{minimize} && 0.5x_1^2 + 0.25x_2^2 \\ &\text{subject to} && e^{x_1} - 2 - x_2 \leq 0 \\ &&& (x_1 - 1)^2 + x_2 - 3 \leq 0 \\ &&& x_1 \geq 0 \\ &&& x_2 \geq 1 \end{aligned}$$



**Contour plot has curves
(no longer lines)**

**Feasible set is
no longer a polyhedron**


Integer optimization

It's still nonlinear optimization

minimize $f(x)$
subject to $x \in \mathbf{Z}$



minimize $f(x)$
subject to $\sin(\pi x) = 0$



**We cannot solve most nonlinear
optimization problems**

Examples of (solvable) nonlinear optimization

Regression

Fit affine function $f(z) = \alpha + \beta z$ to m points (z_i, y_i)

Approximation problem $Ax \approx b$ where

$$A = \begin{bmatrix} 1 & z_1 \\ \vdots & \vdots \\ 1 & z_m \end{bmatrix}, \quad x = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}, \quad b = \begin{bmatrix} y_1 \\ \vdots \\ y_m \end{bmatrix}$$

Goal

$$\text{minimize } \|Ax - b\|$$

1-norm or ∞ -norm \implies **linear optimization**


2-norm \implies **least-squares**

$$\|Ax - b\|_2^2 = \sum_i (f(z_i) - y_i)^2$$

Sparse regression

Regressor selection

minimize $\|Ax - b\|_2^2$
subject to $\text{card}(x) \leq k$
|
(very hard)



Add regularization to the objective

Regularized regression (ridge)

minimize $\|Ax - b\|_2^2 + \gamma\|x\|_2^2$

Regularized regression (lasso)

minimize $\|Ax - b\|_2^2 + \gamma\|x\|_1$

Lasso vs ridge regression



Regularized regression (ridge)

minimize $\|Ax - b\|_2^2 + \gamma\|x\|_2^2$

Regularized regression (lasso)

minimize $\|Ax - b\|_2^2 + \gamma\|x\|_1$

Regularization paths




Portfolio optimization

We have a total of n assets

x_i is fraction of money invested in asset i → **Returns**
 p_i is the relative price change of asset i $p^T x$

p random variable: mean μ , covariance Σ



Convex analysis review

Extended real-value functions

$f(x)$ on $\text{dom } f$

Extended-value extension

$$\tilde{f}(x) = \begin{cases} f(x) & x \in \text{dom } f \\ \infty & x \notin \text{dom } f \end{cases}$$

Always possible to evaluate functions

$$\text{dom } \tilde{f} = \{x \mid \tilde{f}(x) < \infty\}$$

Indicator functions

Indicator function

$$\mathcal{I}_C(x) = \begin{cases} 0 & x \in C \\ \infty & x \notin C \end{cases}$$

Constrained form

minimize $f(x)$
subject to $x \in C$



Unconstrained form


minimize $f(x) + \mathcal{I}_C(x)$

Convex set

Definition

For any $x, y \in C$ and any $\alpha \in [0, 1]$


$$\alpha x + (1 - \alpha)y \in C$$



Convex

Examples

- \mathbf{R}^n
- Hyperplanes
- Hyperspheres
- Polyhedra



Not convex

Examples

- Cardinality constraint $\text{card}(x) \leq k$
- \mathbf{Z}^n
- Any disjoint set



Convex combinations

Convex combination

$\alpha_1 x_1 + \cdots + \alpha_k x_k$ for any x_1, \dots, x_k and $\alpha_1, \dots, \alpha_k$ such that $\alpha_i \geq 0$, $\sum_{i=1}^k \alpha_i = 1$

Convex hull

$$\text{conv } C = \left\{ \sum_{i=1}^k \alpha_i x_i \mid x_i \in C, \alpha_i \geq 0, i = 1, \dots, k, \mathbf{1}^T \alpha = 1 \right\}$$




Cones

Cone

$$x \in C \implies tx \in C \text{ for all } t \geq 0$$

Convex cone

$$x_1, x_2 \in C \implies t_1x_1 + t_2x_2 \in C \text{ for all } t_1, t_2 \geq 0$$





Conic combinations

Conic combination

$\alpha_1 x_1 + \cdots + \alpha_k x_k$ for any x_1, \dots, x_k and $\alpha_1, \dots, \alpha_k$ such that $\alpha_i \geq 0$

Conic hull

$$\left\{ \sum_{i=1}^k \alpha_i x_i \mid x_i \in C, \alpha_i \geq 0, i = 1, \dots, k \right\}$$



Cones

Examples

Nonnegative orthant

$$\mathbf{R}_+^n = \{x \in \mathbf{R}^n \mid x \geq 0\}$$

Norm-cone

$$\{(x, t) \mid \|x\| \leq t\} \quad (\text{if 2-norm, second-order cone})$$


Positive semidefinite cone

$$\mathbf{S}_+^n = \{X \in \mathbf{S}^n \mid z^T X z \geq 0, \quad \text{for all } z \in \mathbf{R}^n\}$$

Normal cone

For any set C and point $x \in C$, we define

$$\mathcal{N}_C(x) = \{g \mid g^T(y - x) \leq 0, \text{ for all } y \in C\}$$



$\mathcal{N}_C(x)$ is **always convex**

What if $x \in \text{int}S$?

Gradient

Derivative

If $f(x) : \mathbf{R}^n \rightarrow \mathbf{R}^m$ continuously differentiable, we define

$$Df(x)_{ij} = \frac{\partial f_i(x)}{\partial x_j}, \quad i = 1, \dots, m, \quad j = 1, \dots, n$$

Gradient

If $f : \mathbf{R}^n \rightarrow \mathbf{R}$, we define

$$\nabla f(x) = Df(x)^T$$

Example

$$f(x) = (1/2)x^T P x + q^T x$$

$$\nabla f(x) = P x + q$$

First-order approximation

$$f(y) \approx f(x) + \nabla f(x)^T (y - x)$$

(affine function of y)

Hessian

Hessian matrix (second derivative)

If $f(x) : \mathbf{R}^n \rightarrow \mathbf{R}$ second-order differentiable, we define

$$\nabla^2 f(x)_{ij} = \frac{\partial^2 f(x)}{\partial x_i \partial x_j}, \quad i, j = 1, \dots, n$$

Example

$$f(x) = (1/2)x^T P x + q^T x$$

$$\nabla^2 f(x) = P$$

Second-order approximation

$$f(y) \approx f(x) + \nabla f(x)^T (y - x) + (1/2)(y - x)^T \nabla^2 f(x)(y - x)$$


(quadratic function of y)

Convex optimization

Convex functions

Convex function

For every $x, y \in \mathbf{R}^n$, $\alpha \in [0, 1]$ $f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y)$



Concave function

f is concave if and only if $-f$ is convex


Convex conditions

First-order

Let f be a continuous differentiable function, then it is convex if and only if $\text{dom} f$ is convex and

$$f(y) \geq f(x) + \nabla f(x)^T (y - x)$$

for all $x, y \in \text{dom} f$



Second-order

If f is twice differentiable, then f is convex if and only if $\text{dom} f$ is convex and

$$\nabla^2 f(x) \succeq 0$$

for all $x \in \text{dom} f$

Verifying convexity

Basic definition (inequality)

First and second order conditions (gradient, hessian)



Hard!

Convex calculus (directly construct convex functions)

- Library of basic functions that are convex/concave
- Calculus rules or transformations that preserve convexity



Easy!

Disciplined Convex Programming

Convexity by construction

General composition rule

$h(f_1(x), f_2(x), \dots, f_k(x))$ is convex when h is convex and for each i

- h is nondecreasing in argument i and f_i is convex, or
- h is nonincreasing in argument i and f_i is concave, or
- f_i is affine

**Only sufficient
condition**

Check your functions at <https://dcp.stanford.edu/>

More details and examples in ORF523

Convex optimization problems

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & g_i(x) \leq 0, \quad i = 1, \dots, m \end{array}$$

$$\begin{array}{ll} f : \mathbf{R}^n \rightarrow \mathbf{R} & \text{Convex objective function} \\ g_i : \mathbf{R}^n \rightarrow \mathbf{R} & \text{Convex constraints functions} \end{array}$$

Convex feasible set

$$C = \{x \mid g_i(x) \leq 0, \quad i = 1, \dots, m\}$$

Modelling software for convex optimization

Modelling tools simplify the formulation of convex optimization problems

- **Construct problems** using library of basic functions
- **Verify convexity** by general composition rule
- Express the problem in input format required by a specific solver

Examples

- CVX, YALMIP (Matlab)
- CVXPY (Python)
- Convex.jl (Julia)

Solving convex optimization problems

CVXPY

$$\begin{aligned} & \text{minimize} && \|Ax - b\|_2 \\ & \text{subject to} && 0 \leq x \leq 1 \end{aligned}$$

```
x = cp.Variable(n)
objective = cp.Minimize(cp.norm(A*x - b))
constraints = [0 <= x, x <= 1]
problem = cp.Problem(objective, constraints)

# The optimal objective value is returned by `problem.solve()`.
result = problem.solve()

# The optimal value for x is stored in `x.value`.
print(x.value)
```

Local vs global minima (optimizers)

$$\begin{aligned} &\text{minimize} && f(x) \\ &\text{subject to} && x \in C \end{aligned}$$

Local optimizer x

$$f(y) \geq f(x), \quad \forall y$$

such that $\|x - y\|_2 \leq R$




Global optimizer x

$$f(y) \geq f(x), \quad \forall y \in C$$

Optimality and convexity

Theorem

For a convex optimization problem, any **local minimum** is a **global minimum**



Optimality and convexity

Proof (contradiction)


Suppose that f is convex and x is a local (not global) minimum for f , i.e.,

$$f(y) \geq f(x), \quad \forall y \text{ such that } \|x - y\|_2 \leq R.$$

Therefore, there exists a feasible z such that $\|z - x\| > R$ and $f(z) < f(x)$.

Consider $y = (1 - \alpha)x + \alpha z$ with $\alpha = \frac{R}{2\|z - x\|_2}$.

Then, $\|y - x\|_2 = \alpha\|z - x\|_2 = R/2 < R$, and by convexity of the feasible set, y is feasible.



By convexity of f we have $f(y) \leq (1 - \alpha)f(x) + \alpha f(z) < f(x)$, which contradicts the local optimum definition.

Therefore, x is globally optimal.



"...in fact, the great watershed in optimization isn't between linearity and nonlinearity, but convexity and nonconvexity."

R. Tyrrell Rockafellar, in SIAM Review, 1993

Nonlinear optimization

Topics of this part of the course

Conditions to characterize minima

Algorithms to find (local) minima

(if applied to **convex problems**, they find **global minima**)

Introduction to nonlinear optimization

Today, we learned to:

- **Define** nonlinear optimization problems
- **Understand** convex analysis fundamentals (sets, cones, functions, and gradients)
- **Verify** convexity and **construct** convex optimization problems
- **Define** convex optimization problems in CVXPY
- **Understand** the importance of *convexity vs nonconvexity* in optimization

Next lecture

- Optimality conditions in nonlinear optimization